


WHAT IS CLAIMED IS:

1. A compound of the formula I



I

wherein

10 G is a C₃ to C₅ branched or straight carbon chain and
 (-O₂CR'), (-OH) and (-O₂C(CH₂)_pCH₃) are attached to any
 available carbon atom along G;

m is an integer from 1 to 4;


n is an integer from 0 to 3;

p an integer from 0 to 16;

15 q is an integer from 0 to 3;


where the sum of m, n and q is 3 or 4; and

-O₂CR' is a fragment of a compound of formula



20 wherein

W is a bicyclic hetroaryl of the structure




25

X is -O-, -S-, -SO₂-, -CHR⁵-, -CHR⁵O-, -CHR⁵S-,
 -CHR⁵SO₂-, -CHR⁵CO- or -CH₂CHR⁵-;

Y is a bond or -CHR⁶-;

Z is an aryl or heteroaryl group of the following structure:



5

A is -CH- or -N-;

B is -O- or -S-;

R¹ is hydrogen, alkyl, aryl or alkenyl;

R² is hydrogen, alkyl, aryl, arylalkyl,
10 heteroarylalkyl or alkenyl;

R³ and R⁴ are each independently hydrogen, halo, trifluoromethyl, cyano, alkyl or alkoxy;


R⁵ and R⁶ are each independently hydrogen, alkyl, aryl, alkenyl, CN, CN₄R^{9A} (tetrazole), CO₂R^{9A}, CONR^{9A}R^{9B} or
15 CONR^{9A}OR^{9B};

R⁷ and R⁸ are each independently hydrogen, halo, trifluoromethyl, cyano, hydroxy, a hydrogen bonding group, alkyl, alkoxy, aryl, arylalkyl, heteroarylalkyl, aryloxy or alkenyl; and

20 R^{9A} and R^{9B} are independently hydrogen, alkyl, arylalkyl, heteroarylalkyl or aryl, or R^{9A} and R^{9B} may optionally be cyclized together to form a ring, wherein said ring may further be substituted with one to three additional hydrogen bonding groups;

25 wherein when R¹, R², R⁵, R⁶, R⁷ and R⁸ are alkyl, aryl, alkenyl, arylalkyl, heteroarylalkyl, alkoxy or aryloxy, R¹, R², R⁵, R⁶, R⁷ and R⁸ may each independently be substituted with 1 to 3 hydrogen bonding groups.

2. The compounds as defined in claim 1 wherein Z is an aryl or heteroaryl group of the structure



5

3. The compound as defined in claim 1 wherein said hydrogen bonding group is selected from the group consisting of OR^{9A} , OCO_2R^{10} , $CONR^{9A}R^{9B}$, CN , NO_2 , CN_4R^{9A} (tetrazole), $COCF_3$, COR^{9A} , CO_2R^{9A} , $CONR^{9A}R^{9B}$, $CONR^{9A}OR^{9B}$,
 10 $C(NR^{9A})NR^{9B}R^{9C}$, $CONR^{9A}SO_2R^{9B}$, SOR^{10} , SO_2R^{10} , SO_3H , $SO_2NR^{9A}R^{9B}$, $SO_2NR^{9A}COR^{9B}$, $SO_2NR^{9A}CONR^{9B}R^{9C}$, $POR^{9A}R^{9B}$, $PO_2R^{9A}R^{9B}$, $PO_3R^{9A}R^{9B}$, $PO_2R^{9A}NR^{9B}R^{9C}$, $NR^{9A}R^{9B}$, $NR^{9A}COR^{9B}$, $NR^{9A}C(NR^{9B})R^{9C}$, $NR^{9A}CO_2R^{9B}$, $NR^{9A}CONR^{9B}R^{9C}$, $NR^{9A}C(NR^{9B})NR^{9C}R^{9D}$, $NR^{9A}SO_2R^{9B}$, $NR^{9A}CONR^{9B}SO_2R^{9C}$, $NR^{9A}SO_2NR^{9B}R^{9C}$, $NR^{9A}POR^{9B}R^{9C}$, $NR^{9A}PO_2R^{9B}R^{9C}$, $NR^{9A}PO_3R^{9B}R^{9C}$ and
 15 $NR^{9A}PO_2R^{9B}NR^{9C}R^{9D}$; wherein

R^{9C} and R^{9D} are each independently hydrogen, alkyl, arylalkyl, heteroarylalkyl or aryl; and


R^{10} is independently alkyl, arylalkyl, heteroarylalkyl, or aryl;

20 wherein R^{9A} , R^{9B} , R^{9C} , R^{9D} or R^{10} may further be substituted with one to three additional hydrogen bonding groups; and wherein two of R^{9A} , R^{9B} , R^{9C} or R^{9D} within the same hydrogen bonding group may optionally be cyclized together to form a ring, wherein said ring may further be
 25 substituted with one to three additional hydrogen bonding groups.


4. The compound as defined in claim 1 where $(-O_2CR')$ represents a fragment of compounds of formula Ib wherein
 30 R^5 or R^6 is CO_2- , or wherein one or more of R^1 , R^2 , R^5 , R^6 , R^7 and R^8 is alkyl, aryl, alkenyl, arylalkyl, heteroarylalkyl, alkoxy or aryloxy, and one of R^1 , R^2 , R^5 ,

R^6 , R^7 and R^8 is substituted with or contains a fragment of structure CO_2^- .

5. The compound as defined in claim 1 wherein
 5 R^1 is hydrogen;
 Z is




W is




10

6. The compound as defined in claim 1 wherein
 R^1 is hydrogen;
 Z is



15

W is



7. The compound as defined in claim 1 wherein W is 5-chloroindol-2-yl.

5 8. The compound as defined in claim 1 where (-O₂CR') represents a fragment of compounds of formula Ib wherein



X is -CHR⁵-, -CHR⁵O-, -CHR⁵S-, -CHR⁵SO₂-, -CHR⁵CO- or -CH₂CHR⁵-;

10 Y is -CHR⁶-; and

R⁵ or R⁶ is CO₂-.



9. The compound as defined in claim 1 wherein W is

15





n is 0.



20 10. The compound as defined in claim 1 having the structure





or





or



5 or




or




10

or



or



5 11. A pharmaceutical composition comprising a compound as defined in Claim 1 and a pharmaceutically acceptable carrier therefor.

10 12. The pharmaceutical composition of claim 11 further comprising at least one additional therapeutic agent selected from the group consisting of other compounds of formula I, anti-diabetic agents, anti-obesity agents, anti-hypertensive agents, anti-atherosclerotic agents and lipid-lowering agents.

15 13. The pharmaceutical composition of claim 12 comprising a compound of formula I and at least one anti-diabetic agent.

20 14. The pharmaceutical composition of claim 13 wherein the antidiabetic agent is at least one agent selected from the group consisting of a biguanide, a sulfonyl urea, a glucosidase inhibitor, a PPAR-alpha

agonist, a PPAR-gamma agonist, a PPAR alpha/gamma dual agonist, an aP2 inhibitor, an SGLT2 inhibitor, a dipeptidyl peptidase-IV inhibitor, an insulin sensitizer, a thiazolidinedione, a glucagon-like peptide-1 (GLP-1),
5 an aldose reductase inhibitor, a sorbitol dehydrogenase inhibitor, insulin and a meglitinide.

15. The pharmaceutical composition of claim 13 wherein the antidiabetic agent is at least one agent
10 selected from the group consisting of metformin, glyburide, glimepiride, glipyrice, glipizide, chlorpropamide, gliclazide, acarbose, miglitol, pioglitazone, troglitazone, rosiglitazone, insulin, Gl-262570, isaglitazone, JTT-501, NN-2344, L895645, YM-440,
15 R-119702, AJ9677, repaglinide, nateglinide, KAD1129, AR-HO39242, GW-409544, KRP297, AC2993, LY315902 and NVP-DPP-728A.

16. The pharmaceutical composition of claim 12 wherein the anti-obesity agent is at least one agent
20 selected from the group consisting of a beta 3 adrenergic agonist, a lipase inhibitor, a serotonin (and dopamine) reuptake inhibitor, a thyroid receptor beta compound and an anorectic agent.

25
17. The pharmaceutical composition of claim 12 wherein the anti-obesity agent is at least one agent selected from the group consisting of orlistat, ATL-962, AJ9677, L750355, CP331648, sibutramine, topiramate,
30 axokine, dexamphetamine, phentermine, phenylpropanolamine and mazindol.

18. The pharmaceutical composition of claim 12 wherein the lipid lowering agent is at least one agent
35 selected from the group consisting of an MTP inhibitor, cholesterol ester transfer protein, an HMG CoA reductase inhibitor, a squalene synthetase inhibitor, a fibric acid

derivative, a cholesterol absorption inhibitor, an ileal Na+/bile cotransporter inhibitor, a bile acid sequestrant, a nicotinic acid derivative, an upregulator of LDL receptor activity, a lipoxygenase inhibitor and an
5 ACAT inhibitor.

19. The pharmaceutical composition of claim 12 wherein the lipid lowering agent is at least one agent selected from the group consisting of pravastatin,
10 lovastatin, simvastatin, atorvastatin, cerivastatin, fluvastatin, nisvastatin, visastatin, fenofibrate, gemfibrozil, clofibrate, avasimibe, TS-962, MD-700, CP-529414, and/or LY295427.

15 20. The pharmaceutical composition of claim 12 comprising a compound of formula I and at least one anti-hypertensive agent.

21. A method for treating or delaying the
20 progression or onset of diabetes, diabetic retinopathy, diabetic neuropathy, diabetic nephropathy, wound healing, insulin resistance, hyperglycemia, hyperinsulinemia, Syndrome X, diabetic complications, elevated blood levels of free fatty acids or glycerol, hyperlipidemia,
25 dislipidemia, obesity, hypertriglyceridemia, atherosclerosis, glucose intolerance, or hypertension which comprises administering to a mammalian patient in need of treatment a therapeutically effective amount of a compound as defined in claim 1.

30 22. The method according to claim 21 further comprising administering, concurrently or sequentially, a therapeutically effective amount of at least one additional therapeutic agent selected from the group
35 consisting of other compounds of formula I, anti-diabetic agents, anti-obesity agents, anti-hypertensive agents, anti-atherosclerotic agents and lipid-lowering agents.

23. A method of inhibiting the enzyme glycogen
phosphorylase which comprises administering to a
mammalian patient in need of treatment a therapeutically
5 effective amount of a compound as defined in claim 1.